



Source of mercury in mining communities of Guyana.

Richard Couture.Natural Resources Canada ricoutur@nrcan.gc.ca
J.D.Lambert.Fisheries and Oceans Canada lambertjd@dfo-mpo.gc.ca

August 21 2003. GGMC Mining Week. Le Méridien Pegasus, Georgetown, Guyana

Background

- Modern gold rush started in the late 1970's;
- Mercury release estimates by Malm (1998), Pfeiffer and Lacerda (1988) and Pfeiffer et al. (1993);
- Environmental consequences of Hg; amalgamation mining on the tropical ecosystems of South America began in the mid- to late-1980s (Pfeiffer and Lacerda, 1988; Martinelli et al., 1988; Lacerda et al., 1989, 1990; 1991a, 1991b; Malm et al. 1990; Pfeiffer et al., 1989, 1991; Lacerda and Salomons, 1992; Nriagu et al., 1992);
- Other source of Hg to the aquatic environment on a regional scale:
 1. Hg concentrations in no known mining areas exhibit values on a regional scale that are comparable to those in basins with extensive mining operations (Forsberg et al., 1995; Malm, 1998; Roulet et al. 1998);
 2. Hg concentrations do not systematically decrease downstream of the mining camps as expected from point sources of contamination (Lechler et al., 2000; Roulet et al., 1998);

Objectives of the study

1. Provide measurements of mercury from amalgam mining areas and from areas of no-known modern mining activity;
2. Investigate if mercury manipulation during the gold amalgamation process can be link to the mercury found in the aquatic environment.

Sampling programme



- Methyl-mercury in:
 - Fish (80 samples)
- Total-mercury in:
 - Water (350 samples)
 - Riverbed sediments (400)
 - Land sediments (100)
 - Fish (1 200)



Results – Objective 1

- **Hg in water:**



- Typical water column vertical distribution shows Hg peak concentration at 1m depth in all mining and non-mining areas;
- The similar trend between turbidity and Hg in unfiltered water suggests equivalent **[Hg]/sediment load** for mining and non-mining areas.



Results – Objective 1

- **Hg in riverbed sediments:**


- 90% of Hg is associated with mud;
- Low [Hg] variability in the mud fraction between mining or non-mining areas;
- Thin mud layer on the riverbed which is more extended downstream mining creeks than upstream and than in non-mining areas.




Results – Objective 1

● Hg in land sediments:

- 90% Hg is associated with mud;
- Low [Hg] variability in the mud fraction between mining or non-mining areas;
- Peak mud abundance found in the overburden;
- Comparable profiles in pristine and mining areas.





Vertical profiles of land sediments



Results – Objective 1

- **Hg in fish:**


- 90% T-Hg is in the form of Me-Hg;
- Mining areas show highest Hg levels in carnivorous fish;
- Omnivorous and herbivorous fish are below 0,5µg Hg/g (WHO) in mining and non-mining areas.



Results – Objective 2

● Source of Hg



- Irrespective of the sampling area (mining or non-mining), [Hg] is related to size class of substrat;
- Irrespective of the sampling area (mining or non-mining) [Hg] is the same on land and on riverbeds;
- As showed from the Potaro river riverbed sediment study, the extension of the mud layer on the riverbed is greater downstream mining creeks.



Results – Objective 2

● Source of Hg



- On land sediments, average concentration of Hg from all size classes is not mining related.
- On riverbed sediments, average Hg concentration for all size classes shows that mining areas contain more Hg (mud) than from non-mining areas;



Results – Objective 2

● Source of Hg

- On the riverbed of the Potaro River and in the White Hole mine facing Region 9), the most abundant fraction of Hg is associated with humic organic matter;
- In the White Hole mine facing, this fraction is located in the overburden.



Conclusion

- Objective 1:

- Hg is trapped in the mud fraction over all land areas but on the riverbeds it is concentrated downstream from mining activities;
- Me-Hg exceeds WHO limits only in carnivorous fish and mainly in mining areas.

- Objective 2:

- No indications that modern amalgam processing is responsible for the concentration of mercury measured in the aquatic environment;
- Strong indications that jetting from land dredges in gold and diamond operations are responsible for the flushing of pre-modern mining mercury from the land overburden to the aquatic environment.